



CAN STATURE, ABDOMINAL PERIMETER AND BMI INDEX PREDICT POSSIBLE CARDIOMETABOLIC RISKS IN FUTURE OBESITY?

Pode índice com estatura, perímetro abdominal e IMC prever possíveis riscos cardiometabólicos em futura obesidade?

Ricardo Wallace das Chagas **LUCAS**^{1,2,3}, Paulo Afonso Nunes **NASSIF**^{1,2,3}, Fernando Issamu **TABUSHI**¹, Denise Serpa Bopp **NASSIF**^{1,2}, Bruno Luiz **ARIEDE**¹, Jose **BRITES-NETO**¹, Osvaldo **MALAFIA**^{1,3}

ABSTRACT - Background: Obesity changes the anatomy of the patient. In addition to the aesthetic change, the high percentage of fat determines evident functional changes. Anthropometric normality in measuring abdominal circumference and height can serve as a basis for measuring cardiometabolic risks of obesity. **Aim:** To verify if it is possible to determine parameters of normality between waist and height in people with normal BMI and fat percentages, to serve as a basis for assessing risks for obesity comorbidities. **Methods:** A sample of 454 individuals with BMI and percentages of fat considered within the normal range was extracted. It was divided into age groups for both men and women between 18 and 25; 26 to 35; 36 to 45; 46 to 55; 56 to 65. A total of 249 men and 205 women were included. **Results:** Regarding the percentage of height as a measure of the abdominal perimeter, the total female sample had an average of $44.2 \pm 1.1\%$ and the male $45.3\% \pm 1.5$. For women, this percentage determined the equation of the waist-height ratio represented by $X = (\text{age} + 217) / 5.875$, and for men $X = (\text{age} + 190.89) / 5.2222$. "X" represents the percentage of the height measurement so that the individual falls into the category of adequate percentage of fat and BMI. **Conclusion:** Between the stature of adult men and women with normal fat percentage and BMI, there is a common numerical relationship, with is on average 44% for women and 45% for men.

HEADINGS: Obesity. Obesidade abdominal. Metabolic syndrome.

RESUMO - Racional: A obesidade modifica a anatomia de seu portador. Além da mudança estética o alto percentual de gordura determina evidentes alterações funcionais. A normalidade antropométrica na mensuração do perímetro abdominal e estatura pode servir de base na mensuração dos riscos cardiometabólicos da obesidade. **Objetivo:** Verificar se é possível determinar parâmetros de normalidade entre cintura e estatura em pessoas que apresentem IMC e percentuais de gordura normais, para servir de base na avaliação de riscos para co-morbidades da obesidade. **Métodos:** Foi extraída amostra de 454 indivíduos com IMC e percentuais de gordura considerados dentro da normalidade. Ela foi dividida em faixas de idade tanto para homens como para mulheres entre 18 a 25; 26 a 35; 36 a 45; 46 a 55; 56 a 65. Totalizou 249 homens e 205 mulheres. **Resultados:** Em relação ao percentual da estatura como medida do perímetro abdominal, a amostra total feminina apresentou média de $44,2 \pm 1,1\%$ e a masculina $45,3\% \pm 1,5$. Para as mulheres este percentual determinou a equação da relação cintura-estatura representado por $X = (\text{idade} + 217) / 5,875$, e para homens $X = (\text{idade} + 190,89) / 5,2222$. "X" representa o percentual da medida da estatura para que o indivíduo se enquadre na categoria de adequados percentual de gordura e IMC. **Conclusão:** Entre a estatura de homens e mulheres adultos possuidores de percentuais de gordura e IMC normais, existe relação numérica comum, sendo em média 44% para mulheres e 45% para homens.

DESCRIPTORIOS: Obesidade. Obesidade abdominal. Síndrome metabólica.

Age range	Percentual relation	
	Women	Men
18 a 25	42.6	43.3
26 a 35	43.8	44.2
36 a 45	44.9	45.8
46 a 55	44.1	46.1
56 a 65	45.4	47.1

Waist / Height normality

Central message

Correlating height, abdominal waist and BMI to create a percentile that defines the normality of this correlation, can contribute to the daily clinical practice of health professionals, as it offers an index of easy interpretation and usability to raise awareness or prevent cardiovascular risk throughout the time. The normal correlation for women is 44% and for men 45%.

Perspective

In recent years, clinical studies have made it clear that the type of body fat distribution can help stratify future cardiovascular risk, and procedures for measuring this distribution can be used on a daily basis. Therefore, correlating height, abdominal waist and BMI to create a percentile that defines the normality of this correlation, may contribute to the daily clinical practice of health professionals, as it indicates an index of easy interpretation and usability to raise awareness or prevent this risk over time. This article details how to reach and use that index.



www.facebook.com/abcdrevista



www.instagram.com/abcdrevista



www.twitter.com/abcdrevista

From the ¹Programa de Pós-Graduação em Princípios da Cirurgia, Faculdade Evangélica Mackenzie do Paraná/Instituto de Pesquisas Médicas, Curitiba, PR, Brasil; ²Instituto Paulo Nassif, Curitiba, PR, Brasil; ³Serviço de Cirurgia Bariátrica e Metabólica, Hospital Universitário Evangélico Mackenzie, Curitiba, PR, Brasil (¹Postgraduate Program in Principles of Surgery, Mackenzie Evangelical Faculty of Paraná/Medical Research Institute, Curitiba, PR, Brazil; ²Paulo Nassif Institute, Curitiba, PR, Brazil; ³Bariatric and Metabolic Surgery Service, University Evangelical Mackenzie Hospital, Curitiba, PR, Brazil).

How to cite this article: Lucas RWC, Nassif PAN, Tabushi FI, Nassif DSB, Ariede BA, Brites-Neto J, Malafía O. Pode índice com estatura, perímetro abdominal e imc prever possíveis riscos cardiometabólicos em futura obesidade? ABCD Arq Bras Cir Dig. 2020;33(2):e1529. DOI: /10.1590/0102-672020190001e1529

Correspondence:
Paulo Afonso Nunes Nassif
Email: paulonassif@terra.com.br

Financial source: This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001
Conflict of interest: none
Received for publication: 06/02/2020
Accepted for publication: 29/05/2020



INTRODUCTION

Obesity has become a worldwide epidemic^{7,27}. Although there are indications that it is metabolically healthy, there is consensus on its harm and various risks present in obese individuals who have visceral fat deposition^{5,20,23,28}. The greater the volume and the percentage of body fat with the increase in body mass index (BMI), there is also deposition in visceral fat³. However, the methodological modalities used to measure total body fat are not directly related to visceral fat, which can be detected by ultrasound, tomography and magnetic resonance.

Despite not having high accuracy, anthropometric measurements have been the most used to assess visceral fat. Among them have been those that analyze the perimeters. The use of the waist/hip ratio and the abdominal waist has shown adequate correlations with visceral fat, when estimated by tomography, and adequately predict cardiometabolic risk⁷.

Regarding the measurement of the abdominal perimeter (waist), the adequacy of the cutoff point established at 102 cm for men and 88 cm for women¹² is questioned for populations of different ethnicities. Some studies with lower levels - 94 cm for men and 80 cm for women - have been considered more appropriate^{6,10,18,20,28}. It is still mentioned that different ethnicities have different somatotypes, and consequently different fat distributions, so that cut-off values predictive of risk in a given population may not be valid for others. The use of height, as it is relatively immutable after adulthood, has served as a basis for the composition of measures more applicable to different populations²¹.

In this sense, an adult individual with normal body composition, regardless of gender and age group, presents body measurements with numerical values related to height.

Thus, the aim of this study was to study the percentage waist-height ratio in patients with normal fat percentages, and to format a table (Table 3) relating it to age, in order to propose parameters for the assessment of cardiometabolic risks in future obesity.

METHODS

This prospective descriptive study correlated data collected from the Bariatric and Metabolic Surgery Service of Hospital Universitário Evangélico Mackenzie in Curitiba, PR, Brazil. The sample initially had 1,717 individuals. The data of interest for the study were: height, total body mass (weight), the percentage of fat and the smallest abdominal circumference. The variables were collected as follows: height using a portable stadiometer with a 1 mm metric scale; total body mass by a calibrated portable digital scale with a capacity of 150 kg and a margin of 100 g; abdominal perimeter, with an inextensible measuring tape and 0.1 cm margin; fat percentage, by Omron bipolar bioimpedance equipment, model HBF-306.

With body composition considered adequate within the initial sample, 454 adult individuals of both genders were extracted in the age group between 18 and 65 years old and who had a normal percentage of fat and BMI (Pollock and Wilmore, 1993). They were divided into groups by age groups, namely: 18 to 25 (n=165 ♂ 81♀); 26 to 35 (n=53♂80♀); 36 to 45 (n=15♂25♀); 46 to 55 (n=13♂ 13♀); 56 to 65 (n=3♂ 6♀, Figure 1). There were 249 men and 205 women.

PERCENTUAL DE GORDURA X IMC (HOMENS)						
Nível /idade		18 - 25	26 - 35	36 - 45	46 - 55	56 - 65
Atletico/Risco	≦ou <12 kg/m ²	(*) 4 a 7 %	(*) 8 a 11%	(*) 10 a 15%	(*) 12 a 17%	(*) 13 a 19%
Atletico	12.1 a 18.5 kg/m ²	8 a 11%	12 a 15%	16 a 18%	18 a 20%	20 a 21%
Ótimo	18.6 a 19.9 kg/m ²	12 a 13%	16 a 18%	19 a 21%	21 a 23%	22 a 23%
Saudável	20 a 24.9 kg/m ²	14 a 16%	19 a 21%	22 a 24%	24 a 25%	24 a 25%
Sobrepeso	25 a 29.9 kg/m ²	17 a 20%	22 a 24%	25 a 27%	26 a 27%	26 a 27%
Obeso Grau I	30 a 34.9 kg/m ²	21 a 24%	25 a 27%	28 a 30%	28 a 31%	28 a 31%
Obeso Grau II	35 a 39.9 kg/m ²	25 a 36%	28 a 36%	31 a 39%	32 a 38%	32 a 38%

PERCENTUAL DE GORDURA X IMC (MULHERES)						
Nível /idade		18 - 25	26 - 35	36 - 45	46 - 55	56 - 65
Atletico/Risco	≦ou <12 kg/m ²	(*) 13 a 16 %	(*) 14 a 17%	(*) 16 a 19%	(*) 17 a 21%	(*) 18 a 22%
Atletico	12.1 a 18.5 kg/m ²	17 a 19%	18 a 20%	20 a 23%	22 a 25%	24 a 26%
Ótimo	18.6 a 19.9 kg/m ²	20 a 22%	21 a 23%	24 a 26%	26 a 28%	27 a 29%
Saudável	20 a 24.9 kg/m ²	23 a 25%	24 a 26%	27 a 29%	29 a 31%	30 a 32%
Sobrepeso	25 a 29.9 kg/m ²	26 a 28%	27 a 29%	30 a 32%	32 a 34%	33 a 35%
Obeso Grau I	30 a 34.9 kg/m ²	29 a 31%	30 a 33%	33 a 36%	35 a 38%	36 a 38%
Obeso Grau II	35 a 39.9 kg/m ²	32 a 43%	34 a 49%	37 a 48%	39 a 50%	39 a 49%

FIGURE 1 - Pollock & Wilmore (1993) table showing BMI vs. % of fat by age group and gender in relation to the state of body complexion

Statistical analysis

For data treatment, the Excel statistical program was used. Statistical procedures were carried out following age, height, abdominal circumference, fat percentage and waist-to-height ratio. Pearson's correlation coefficient (r), mean and standard deviation of all variables were used, and the 95% confidence interval for mean abdominal circumference. The waist-to-height ratios of the population were transcribed in percentages, and the averages of the abdominal perimeter were determined.

RESULTS

Separating the genders, the standard deviation of the mean of each age group was calculated. Adding all of them, the female sample had an average of 44.2+1.1% in height.

TABLE 1 - Mean and standard deviation of each stratum of female age group by analyzed parameter (n=205)

Age range	<Abdominal perimeter	Height	Percentual relation	BMI	%Fat
18 a 25	70	164.4	42.6	20.4	19.6
26 a 35	71.7	163.8	43.8	20.9	21.8
36 a 45	73.3	163.2	44.9	21.9	25.7
46 a 55	71.2	161.2	44.1	21.5	25.8
56 a 65	71.7	157.8	45.4	21.8	31.3
Mean	71.6	162.1	44.2	21.3	24.8
SD	1.2	2.7	1.1	0.6	4.5

The male sample had an average height of 45.3%.

TABLE 2 - Mean and standard deviation of each stratum of male age group per analyzed parameter (n=249)

Age range	<Abdominal perimeter	Height	Percentual relation	BMI	% Fat
18 a 25	75.7	174.9	43.3	21.7	10.3
26 a 35	77.7	175.8	44.2	22	12.6
36 a 45	80.8	176.3	45.8	23	18
46 a 55	79.9	173.6	46.1	22.9	20.1
56 a 65	83.3	176.7	47.1	21.6	20
Mean	79.5	175.2	45.3	22.2	16.2
SD	2,9	1,2	1,5	0,7	4,5

Taking into account the interval between the smallest and the largest range of each gender, a percentage range was found that varied from 39.5% to 47.9% for women and 40.4% to 49.1% for men.

When using Pearson's correlation between abdominal perimeters and the fat percentage of the female sample, a strong positive correlation was found: R²=0.8559.




FIGURE 2 – Correlation between abdominal circumference and fat percentage in women

There was also a strong positive correlation ($R^2=0.9406$) when using the Pearson correlation method between the abdominal perimeters analyzed and the fat percentage of the male sample.




FIGURE 3 - Correlation between abdominal circumference and fat percentage in men

The sample presented nominally for the values of waist in percentage of the height, interval between 40.4% to 49.1% for men and 39.5% to 47.9% for women. Data divided into age groups are shown in Table 3.

TABLE 3 - Stratum of normal waist-height in relation to gender and age groups in people with normal BMI

Age range	Percentual relation	
	Women	Men
18 a 25	42.6	43.3
26 a 35	43.8	44.2
36 a 45	44.9	45.8
46 a 55	44.1	46.1
56 a 65	45.4	47.1

DISCUSSION

The focus on the correlation between cardiometabolic disorders present in certain populations and anthropometric or body composition signs, characterizes the waist-height advantage over other anthropometric indices. The waist-to-height ratio is a good discriminator of abdominal obesity related to cardiometabolic risk factors^{3,5,13,19,22}.

In a study carried out with a sample of 55,563 adults of both genders in Taiwan, with the objective of identifying the cutoff points of the waist-to-height ratio to discriminate at least one cardiovascular risk factor (diabetes, hypertension or dyslipidemia), values of 0.48 were found and 0.45 for men and women, respectively, that is, 48% and 45% of height for men and women. Still in Taiwan, using 38,556 subjects of both genders as a sample, there was a strong association

between waist-to-height ratio with arterial hypertension, glucose intolerance, diabetes and dyslipidemia²⁴.

Analyzing specifically the results of this research, there was a tendency in the division by strata of age group to increase the percentage of fat with age, even with the individual within the normal BMI range, as shown in Figure 1.

Taking into account that the cardiometabolic risk is higher in older individuals compared to younger ones, there is a correlation with the increase in the measurement of the abdominal perimeter and the percentage of fat found in the research, even in individuals with adequate body composition³.

When applying Pearson's regression equation to the columns referenced by the fat percentage and waist circumference, a strong positive correlation was found for both men and women. This finding corroborates the finding of the waist-to-height ratio when looking for factors associated with central obesity in adults on a population basis²².

The stratum of all age groups of men was consistent with the research by Lucas et al.²⁰ who, by separating men between 18 and 25 years old, demonstrated that the 43% height range was adequate with a very strong positive correlation (Pearson $r=0.778$) between the fat percentage and the abdominal perimeter. This data also coincides with the assertion that the higher the percentage of fat, the greater the waist circumference^{3,13,14}. Likewise, it demonstrated a concern for the division of the waist-to-height ratio by strata, in view of the physiological change in body composition with changes in age groups. The loss of lean mass with increasing age may be related to the decrease in its percentage, but it can also increase the fat mass and consequently the fat percentage. The correlation of this physiological finding cannot be exempted from the analysis of the level of physical activity of the studied population, so that anthropometric measures can make sense²¹.

The advantage for the male sample over the correlation between the percentage of fat and the abdominal perimeter is in line with what refers to another anthropometric method of perimetric relationship, that is, using the waist-hip ratio. This reason for women acquires lower values than for men due to the gender trait, which allows the percentage of fat for women - even the essential one - to be higher than for men⁸. Thus, the finding of an unfavorable correlation for women is understandable, even though Pearson's correlation is strong.

CONCLUSION

It was possible to determine parameters of normality between waist and height in people with BMI and percentages of normal fat, creating indices (Table 3) for different age groups to assist in the clinical evaluation regarding the risks of comorbidities promoted by obesity.

REFERENCES

- Ashwell M, Gibson S. Waist to height ratio is a simple and effective obesity screening tool for cardiovascular risk factors: analysis of data from the British National Diet and Nutrition Survey of adults aged 19–64 years. *Obes Facts* 2009; 2: 97–103.
- Ashwell M, Lejeune S, Mcpherson K. Ratio of waist circumference to height may be better indicator of need for weight management. *BMJ* 1996; 312: 377.
- Ashwell M. Shape: the waist-to-height ratio is a good, simple screening tool for cardiometabolic risk. *Nutr Today* 2011; 46: 85–89.
- Ashwell M. The Ashwell shape chart—a public health approach to the metabolic risks of obesity. *Int J Obes Relat Metab Disord* 1997; 22: S213.
- Barbosa LS, Scala LCN, Ferreira MG. Associação entre marcadores antropométricos de adiposidade corporal e hipertensão arterial na população adulta de Cuiabá, Mato Grosso. *Rev. Bras. Epidemiol.*, v.12, n.2, p. 237–246, 2009.
- Barbosa PJB, et al. Critério de obesidade central em população brasileira: impacto sobre a síndrome metabólica. *Arq. Bras. Cardiol.* v.87 n.4, p.407–414, 2006.

7. Barros F, Negrão MG, Negrão GG. Weight loss comparison after sleeve and roux-en-y gastric bypass: systematic review. *ABCD, arq. bras. cir. dig.*, 2019, vol.32, no.4. ISSN 0102-6720 B.
8. Dalton M, et al. Waist circumference, waist-hip ratio and body mass index and their correlation with cardiovascular disease risk factors in Australian adults. *J Intern. Med.*, v.254, n.6, p.555-63. 2003.
9. Dâmaso A. Obesidade. Rio de Janeiro: Medsi, 2003. 590p.
10. Ferreira SR, Japanese-Brazilian Diabetes Study Group, et al. Prevalence and 7-year incidence of type II diabetes mellitus in a Japanese-Brazilian population: an alarming public health problem. *Diabetologia* 2002;45:1635-8.
11. Ferreira MG, et al. Acurácia da circunferência da cintura e da relação cintura/quadril como preditores de dislipidemias em estudo transversal de doadores de sangue de Cuiabá, Mato Grosso, Brasil. *Cad. Saúde Pública*, v.22, n.2, p.307-314, 2006.
12. Guizoni M, Lucas RWC. Estudo da relação da estatura com a menor circunferência torácico abdominal e o percentual de gordura. Monografia apresentada para conclusão de Aperfeiçoamento/Especialização em Fisioterapia – Faculdade de Educação Física e Fisioterapia de Jacarezinho, 2006.
13. Han TS, Van Leer EM, Seidell JC, Lean MEJ. Waist circumference action levels in the identification of cardiovascular risk factors: prevalence study in a random sample. *BMJ*, n.311, p.1401-5. 1995.
14. Hsieh SD, et al. Health risks among Japanese men with moderate body mass index. *Int J Obes Relat Metab Disord* 2000; 24: 358–362. | Article | PubMed | ChemPort
15. Hsieh SD, et al. Anthropometric obesity indices in relation to age and gender in Japanese adults. *Tohoku J Exp Med* 2000; 191: 79–84.
16. Hsieh SD, et al. Regular physical activity and coronary risk factors in Japanese men. *Circulation* 1998; 97: 661–665. HSIEH SD, YOSHINAGA H. Do people with similar waist circumference share similar health risks irrespective of height? *Tohoku J Exp Med* 1999; 188: 55–60.
17. Huang KC, et al. Four anthropometric indices and cardiovascular risk factors in Taiwan. *Int J Obes Relat Metab Disord* 2002;26(8):1060-8.
18. Lerario DDG, et al. Excesso de peso e gordura abdominal para a síndrome metabólica em nipo-brasileiros. *Rev. Saúde Pública*, v.36, p.4-11, 2002.
19. Little P, Byrne CD. Abdominal obesity and the "hypertriglyceridaemic waist" phenotype. It's probably not yet time to implement screening. *BMJ* 2001; 322: 687–689.
20. Lucas RWC. Análise da relação entre a estatura e o perímetro abdominal em indivíduos portadores de percentuais normais de gordura. Dissertação de Mestrado. UDESC, 2010.
21. Nonino CB, et al. Is there any change in phenotypic characteristics comparing 5 to 10 years of follow-up in obese patients undergoing Roux-en-Y gastric bypass?. *ABCD, arq. bras. cir. dig.*, 2019, vol.32, no.3. ISSN 0102-6720 C.
22. Oh JY, et al. Prevalence and factor analysis of metabolic syndrome in an urban Korean population. *Diabetes Care*, v.27, p.2027-2032, 2004.
23. Pitanga FJG, Lessa I. Associação entre indicadores antropométricos de obesidade e risco coronariano em adultos na cidade de Salvador, Bahia, Brasil. *Rev. Bras. Epidemiol.*, v.10, n.2, p.239-248, 2007.
24. Pitanga FJG, Lessa I. Indicadores antropométricos de obesidade como instrumento de triagem para risco coronariano elevado em adultos na cidade de Salvador – Bahia. *Arq. Bras. Cardiol.*, v.85, n.1, p.26-31, 2005.
25. Pitanga FJG, Lessa I. Razão cintura-estatura como discriminador de risco coronariano de adultos. *Rev. Assoc. Med. Bras.* [online], v.52, n.3, ISSN 0101-4230, 2006.
26. Souza TF, et al. Fatores associados à obesidade central em adultos de Florianópolis, Santa Catarina: estudo de base populacional. *Rev Bras Epidemiol* 2011; 14(2): 296-309.
27. Tarastchuk JCE, et al. Obesidade e intervenção coronariana: devemos continuar valorizando o Índice de Massa Corpórea? *Arq. Bras. Cardiol.* v.90, n.5, p.311-316, 2008.
28. Tonatto-Filho AJ, et al. Bariatric surgery in brazilian public health system: the good, the bad and the ugly, or a long way to go. Yellow sign!. *ABCD, arq. bras. cir. dig.*, 2019, vol.32, no.4. ISSN 0102-6720 A.
29. Zilberstein B, Santo MA, Carvalho MH. Critical Analysis Of Surgical Treatment Techniques Of Morbid Obesity. *ABCD, arq. bras. cir. dig.*, 2019, vol.32, no.3. ISSN 0102-6720 D.